



Opioid Prescribing by Foot and Ankle Surgeons

Approved by the Board of Directors July 2022

The Center for Disease Control and Prevention (CDC) (2018) reports a fivefold increase in opioid deaths since 1999. In addition, the CDC reported (2018) 115 patient deaths daily from opioid overdose in the year 2016. Opioid prescribing has dramatically increased since the 1990s. Both prescription and non-prescription opioid usage has resulted in an opioid crisis in America. Opioid overdose is the leading cause of accidental deaths in young adults, associated with a higher risk of postoperative death, and increases the risk of fall and fractures.

American College of Foot and Ankle Surgeons (ACFAS) members subscribe to the highest ethical, professional, and legal standards and recognize that patients need to have access to appropriate pain relief. ACFAS and its members also recognize the importance of working towards a solution to the opioid crisis.

Opioids are often indicated for the treatment of severe short-lived pain during acute events such as trauma or surgery. Although opioids can be highly addictive, opioid addiction rarely emerges when opioids are used for short-term treatment of pain, except among a few highly susceptible individuals. For the foot and ankle surgeon, the usage of opioids in the acute setting encompasses the vast majority of prescribing. Despite this stated value of opioids, the role of opioids in the treatment of chronic pain has come under scrutiny. Recent open-ended and indiscriminate long-term prescribing of opioids in the United States has led to high rates of prescription opioid misuse and abuse, unacceptable death rates, and an enormous burden to the public. Chronic pain treatment strategies that focus on improving long-term quality of life should include consideration of a referral to a pain management specialist by the foot and ankle surgeon.

Standardized Opioid Protocols for the Foot and Ankle Surgeon

These recommendations are for the benefit of the patient. Foot and ankle surgeons should develop protocols and procedures that should include consideration of:

- Individual state guidelines where available.
- Guidelines for the acute pain patient suffering from trauma.
- Guidelines for the acute pain following surgery.
- Identification of patients at high risk for opioid addiction, substance use disorder, or an opioid-related adverse drug event.
- References related to opioid use and alternative pain management solutions.



- Establishing ranges for acceptable amounts and durations for patients during their post-operative period.
- Utilizing a pain management specialist when appropriate.
- Restricting the usage of opioids beyond the immediate postoperative period.
- Restricting the usage of extended release opioids as most patients are treated for acute pain.
- Limiting opioid prescription size established by your set ranges and acceptable amounts

ACFAS strongly advocates continued research, organizational collaborations, adoption of best practices in opioid prescribing, identify ways to minimize opioid risk and find effective alternatives to opioids for the treatment of various pain problems. ACFAS encourages members to refer to federal and state resources for the latest guidelines.

Bibliography

1. Contextual evidence review for the CDC guideline for prescribing opioids for chronic pain - United States, 2016. CDC Stacks, Public Health Publications, March 18, 2016.
2. Aparasu RR, Chatterjee S. Use of narcotic analgesics associated with increased falls and fractures in elderly patients with osteoarthritis. *Evid Based Med*. 2014 Feb;19(1):37-8. doi: 10.1136/eb-2013-101401. Epub 2013 Aug 13. PubMed PMID:23942986.
3. Aparasu RR, Chatterjee S. <http://www.ncbi.nlm.nih.gov/pubmed/23942986>
4. Attal N, Cruccu G, Baron R, Haanpaa M, Hansson P, Jensen TS, Nurmikko T. EFNS guidelines on the pharmacological treatment of neuropathic pain: 2010 revision. *Eur J Neurol* 2010;17(9):113-e88.
5. Baron MJ, McDonald PW. Significant pain reduction in chronic pain patients after detoxification from high-dose opioids. *J Opioid Manag* 2006;2(5):277-82.
6. Boscarino JA, Rukstalis MR, Hoffman SN, Han JJ, Erlich PM, Ross S, Gerhard GS, Stewart WF. Prevalence of prescription opioid-use disorder among chronic pain patients: comparison of the DSM-5 vs. DSM-4 diagnostic criteria. *J Addict Dis* 2011;30(3):185-94.
7. Carey B. Prescription Painkillers Seen as a Gateway to Heroin. *The New York Times*, Feb. 11, 2014. <https://www.nytimes.com/2014/02/11/health/prescription-painkillers-seen-as-a-gateway-to-heroin.html>
8. Case A, Deaton A. Rising morbidity and mortality in midlife among white non-Hispanic Americans in the 21st century. *Proc Natl Acad Sci USA* 2015; 112(49):15078-83.
9. Chou R, Deyo R, Devine B, Hansen RL, Sullivan S, Jarvik JG, Blazina I, Dana T, Bougatsos C, Turner J. The effectiveness and risks of long-term opioid treatment of chronic pain. Rockville (MD): Agency for Healthcare Research and Quality (US); 2014 Sep. (Evidence Reports/Technology Assessments, No. 218). Report No.: 14-E005-EF.
10. Cunningham JL, Evans MM, King SM, Gehin JM, Loukianova LL. Opioid Tapering in Fibromyalgia Patients: Experience from an Interdisciplinary Pain Rehabilitation Program. *Pain Med* 2016;17(9):1676-85.
11. Dillie KS, Fleming MF, Mundt MP, French MT. Quality of life associated with daily opioid therapy in a primary care chronic pain sample. *J Am Board Fam Med* 2008;21(2):108-17.



12. Dunn KM, Saunders KW, Rutter CM, Banta-Green CJ, Merrill JO, Sullivan MD, Weisner CM, Silverberg MJ, Campbell CI, Psaty BM, Von Korff M. Opioid prescriptions for chronic pain and overdose: a cohort study. *Ann Intern Med* 2010;152(2): 85-92.
13. Eriksen J, Sjogren P, Bruera E, Ekholm O, Rasmussen NK. Critical issues on opioids in chronic non-cancer pain. An epidemiological study. *Pain* 2006;125:172-9.
14. Finlayson RE, Maruta T, Morse RM, Martin MA. Substance dependence and chronic pain: experience with treatment and follow-up results. *Pain* 1986;26(2):175-80.
15. Frank JW, Lovejoy TI, Becker WC, Morasco BJ, Koenig CJ, Hoffecker L, Dischinger HR, Dobscha SK, Krebs EE. Patient Outcomes in Dose Reduction or Discontinuation of Long-Term Opioid Therapy: A Systematic Review. *Ann Intern Med* 2017;167(3):181-91.
16. Goldenberg DL, Clauw DJ, Palmer RE, Clair AG. Opioid Use in Fibromyalgia: A Cautionary Tale. *Mayo Clin Proc* 2016;91(5):640-8.
17. Gomes T, Mamdani MM, Dhalla IA, Paterson JM, Juurlink DN. Opioid dose and drug- related mortality in patients with nonmalignant pain. *Arch Intern Med* 2011;171(7):686-91.
18. Krebs EE, Gravely A, Nugent S, Jensen AC, DeRonne B, Goldsmith ES, Kroenke K, Bair MJ, Nooraloochi S. Effect of opioid vs nonopioid medications on pain-related function in patients with chronic back pain or hip or knee osteoarthritis pain: The SPACE Randomized Clinical Trial. *JAMA* 2018;319(9):872-82.
19. LeeD, ArmaghaniS, ArcherKR, et al. Preoperative Opioid Use as a Predictor of Adverse Postoperative Self-Reported Outcomes in Patients Undergoing Spine Surgery. *J Bone Joint Surg Am.* 2014 Jun 4;96(11):e89 <http://www.ncbi.nlm.nih.gov/pubmed/24897746>
20. Manchikanti L, Singh A: Therapeutic opioids: A ten-year perspective on the complexities and complications of the escalating use, abuse, and nonmedical use of opioids. *Pain Physician* 2008;11(2 suppl):S63-S88. MedlineWeb of Science <http://www.ncbi.nlm.nih.gov/pubmed/18443641>
21. Noble M, Treadwell JR, Tregear SJ, Coates VH, Wiffen PJ, Akafomo C, Schoelles KM. Long-term opioid management for chronic noncancer pain. *Cochrane Database Syst Rev* 2010(1):
22. Okie S. A flood of opioids, a rising tide of deaths. *N Engl J Med.* 2010 Nov 18;363(21):1981-5. doi: 10.1056/NEJMp1011512. <http://www.nejm.org/doi/full/10.1056/NEJMp1011512> Erratum in: *N Engl J Med.* 2011 Jan 20;364(3):290. PubMed PMID: 21083382.
23. Palmer RE, Carrell DS, Cronkite D, Saunders K, Gross DE, Masters E, Donevan S, Hylan TR, Von Kroff M. The prevalence of problem opioid use in patients receiving chronic opioid therapy: computer-assisted review of electronic health record clinical notes. *Pain* 2015;156(7):1208-14.
24. Sjogren P. Epidemiology of chronic pain and critical issues on opioid use. *Pain* 2011;152(6): 1219-20.
25. Toblin RL, Mack KA, Perveen G, Paulozzi LJ. A population-based survey of chronic pain and its treatment with prescription drugs. *Pain* 2011;152(6):249-55.

The American College of Foot and Ankle Surgeons is a professional society of 7,600 foot and ankle surgeons. Founded in 1942, the College's mission is to promote research and provide continuing education for the foot and ankle surgical specialty, and to educate the general public on foot health and conditions of the foot and ankle through its patient education website, FootHealthFacts.org.